

## Research articles using the CS-1 or earlier equivalent models

- Babes, A., Amuzescu, B., Krause, U., Scholz, A., Flonta, M.L. and Reid, G., Cooling inhibits capsaicin-induced currents in cultured rat dorsal root ganglion neurones, *Neurosci Lett.*, 317 (2002) 131-4.
- Babes, A., Zorzon, D. and Reid, G., Two populations of cold-sensitive neurons in rat dorsal root ganglia and their modulation by nerve growth factor, *Eur J Neurosci.*, 20 (2004) 2276-82.
- Babes, A., Zorzon, D. and Reid, G., A novel type of cold-sensitive neuron in rat dorsal root ganglia with rapid adaptation to cooling stimuli, *Eur J Neurosci.*, 24 (2006) 691-8.
- Cao, E., Cordero-Morales, J.F., Liu, B., Qin, F. and Julius, D., TRPV1 channels are intrinsically heat sensitive and negatively regulated by phosphoinositide lipids, *Neuron*, 77 (2013) 667-79.
- Caspani, O., Zurborg, S., Labuz, D. and Heppenstall, P.A., The contribution of TRPM8 and TRPA1 channels to cold allodynia and neuropathic pain, *PLoS One.*, 4 (2009) e7383.
- Chuang, H.H., Neuhausser, W.M. and Julius, D., The super-cooling agent icilin reveals a mechanism of coincidence detection by a temperature-sensitive TRP channel, *Neuron.*, 43 (2004) 859-69.
- Ciobanu, C., Reid, G. and Babes, A., Acute and chronic effects of neurotrophic factors BDNF and GDNF on responses mediated by thermo-sensitive TRP channels in cultured rat dorsal root ganglion neurons, *Brain Res.*, 1284 (2009) 54-67.
- Daniels, R.L., Takashima, Y. and McKemy, D.D., Activity of the neuronal cold sensor TRPM8 is regulated by phospholipase C via the phospholipid phosphoinositol 4,5-bisphosphate, *J Biol Chem.*, 284 (2009) 1570-82.
- Endres-Becker, J., Heppenstall, P.A., Mousa, S.A., Labuz, D., Oksche, A., Schafer, M., Stein, C. and Zollner, C., Mu-opioid receptor activation modulates transient receptor potential vanilloid 1 (TRPV1) currents in sensory neurons in a model of inflammatory pain, *Mol Pharmacol.*, 71 (2007) 12-8.
- Gracheva, E.O., Ingolia, N.T., Kelly, Y.M., Cordero-Morales, J.F., Hollopeter, G., Chesler, A.T., Sanchez, E.E., Perez, J.C., Weissman, J.S. and Julius, D., Molecular basis of infrared detection by snakes, *Nature*, 464 (2010) 1006-11.
- Gracheva, E.O., Cordero-Morales, J.F., Gonzalez-Carcacia, J.A., Ingolia, N.T., Manno, C., Aranguren, C.I., Weissman, J.S. and Julius, D., Ganglion-specific splicing of TRPV1 underlies infrared sensation in vampire bats, *Nature*, 476 (2011) 88-91.
- Linte, R.M., Ciobanu, C., Reid, G. and Babes, A., Desensitization of cold- and menthol-sensitive rat dorsal root ganglion neurones by inflammatory mediators, *Exp Brain Res.*, 178 (2007) 89-98.
- Myers, B.R., Bohlen, C.J. and Julius, D., A yeast genetic screen reveals a critical role for the pore helix domain in TRP channel gating, *Neuron.*, 58 (2008) 362-73.
- Myers, B.R., Sigal, Y.M. and Julius, D., Evolution of thermal response properties in a cold-activated TRP channel, *PLoS One*, 4 (2009) e5741.
- Neacsu, C., Babes, A., The M-Channel blocker linopirdine is an agonist of the capsaicin receptor TRPV1, *J Pharmacol Sci* 114 (2010) 332 – 40
- Neacsu, C., Ciobanu, C., Barbu, I., Toader, O., Szegli, G., Kerek, F., Babes, A., Substance MCS-18 isolated from *Helleborus purpurascens* is a potent antagonist of the capsaicin receptor, TRPV1, in rat cultured sensory neurons, *Physiol. Res.*, 59 (2010) 289-98.
- Pena, F., Amuzescu, B., Neaga, E., Flonta, M.-L., Thermodynamic properties of hyperpolarization-activated current ( $I_h$ ) in a subgroup of primary sensory neurons, *Exp. Brain Res.* 173 (2006) 282–90.
- Prescott, E.D. and Julius, D., A modular PIP2 binding site as a determinant of capsaicin receptor sensitivity, *Science.*, 300 (2003) 1284-8.
- Reid, G. and Flonta, M.L., Physiology. Cold current in thermoreceptive neurons, *Nature.*, 413

- (2001) 480.
- Reid, G. and Flonta, M., Cold transduction by inhibition of a background potassium conductance in rat primary sensory neurones, *Neurosci Lett.*, 297 (2001) 171-4.
- Reid, G., Babes, A. and Pluteanu, F., A cold- and menthol-activated current in rat dorsal root ganglion neurones: properties and role in cold transduction, *J Physiol.*, 545 (2002) 595-614.
- Reid, G. and Flonta, M.L., Ion channels activated by cold and menthol in cultured rat dorsal root ganglion neurones, *Neurosci Lett.*, 324 (2002) 164-8.
- Selescu, T., Ciobanu, A.C., Dobre, C., Reid, G. and Babes, A., Camphor activates and sensitizes transient receptor potential melastatin 8 (TRPM8) to cooling and icilin, *Chem Senses*, 38 (2013) 563-75.
- Takashima, Y., Daniels, R.L., Knowlton, W., Teng, J., Liman, E.R. and McKemy, D.D., Diversity in the neural circuitry of cold sensing revealed by genetic axonal labeling of transient receptor potential melastatin 8 neurons, *J Neurosci.*, 27 (2007) 14147-57.
- Wang, H., Schupp, M., Zurborg, S. and Heppenstall, P.A., Residues in the pore region of *Drosophila* transient receptor potential A1 dictate sensitivity to thermal stimuli, *J Physiol*, 591 (2013) 185-201.
- Zurborg, S., Yurgionas, B., Jira, J.A., Caspani, O. and Heppenstall, P.A., Direct activation of the ion channel TRPA1 by Ca<sup>2+</sup>, *Nat Neurosci.*, 10 (2007) 277-9.

By journal:

Nature 3  
Neuron 3  
Neurosci Lett 3  
PLOS One 2  
J Physiol 2  
Eur J Neurosci 2  
Exp Brain Res 2  
Science 1  
Nat Neurosci 1  
J Neurosci 1  
J Pharmacol Sci 1  
J Biol Chem 1  
Mol Pharmacol 1  
Physiol Res 1  
Brain Res 1  
Chem Senses 1